International Journal of Enhanced Research in Educational Development (IJERED)
ISSN: 2320-8708, Vol. 14 Issue 5, September-October, 2026

Effects of Temperature Variation on Fish Growth

Dr. Rekha Rani

Assistant Professor, Department of Zoology, Allama Iqbal College, Bihar Sharif, Nalanda, Bihar, India

A Comprehensive Review of Environmental Physiology and Bioenergetics

DOI: https://doi.org/10.55948/IJERED.2026.0927

ABSTRACT

Water temperature serves as the abiotic 'master factor' governing the physiology, metabolic demands, and overall growth performance of teleost fishes. Because fish are ectothermic aquatic organisms, their core metabolic functions, enzyme kinetics, and feed conversion efficiencies are inextricably tied to thermal dynamics. This paper reviews the physiological and bioenergetic mechanisms through which temperature variations dictate somatic fish growth. We analyze the concept of Scope for Growth (SFG), the temperature-dependence of feed intake and gastric emptying times, the metabolic costs associated with sub-optimal thermal ranges, and ecological phenomena such as the Temperature-Size Rule (TSR). Additionally, we examine the compounding effects of climate-driven thermal fluctuations on dissolved oxygen mechanics, cellular stress pathways, and aquaculture bioenergetics. Understanding these thermal constraints is critical for optimizing aquaculture management strategies and predicting wild fish population shifts amid ongoing global environmental changes.

INTRODUCTION

As classic poikilotherms, aquatic ectotherms are fundamentally dependent on ambient water temperature to dictate their internal biological activity. Temperature influences nearly every step of fish physiology—ranging from oxygen diffusion and basal metabolic rate to digestion, hormonal secretion, and cellular growth.

In natural ecosystems and aquaculture environments, temperature is rarely static; diurnal cycles, seasonal turnover, thermal effluents, and global climate shifts expose fish to varying degrees of thermal fluctuation. Somatic growth—defined as the accumulation of muscular, skeletal, and visceral tissue over time—serves as a comprehensive integration of a fish's bioenergetic state. When environmental temperature strays from a species-specific optimal range, energy allocation shifts away from protein synthesis and somatic growth toward homeostatic maintenance and physiological stress mitigation.

Bioenergetic Foundations: Scope for Growth and Metabolic Scaling

The Q10 Temperature Coefficient

At the molecular level, biological reaction rates accelerate with increased thermal energy. The rate of metabolic acceleration within an organism's physiological tolerance zone is often quantified using the Q10 temperature coefficient:

$$Q10 = (R2 / R1) ^ [10 / (T2 - T1)]$$

where R1 and R2 represent metabolic rates measured at temperatures T1 and T2 (in °C), respectively. For most teleost species, the baseline metabolic Q10 value approximates 2.0 to 2.5 within their normal thermal performance curve. This means that a 10°C increase in ambient water temperature roughly doubles basal metabolic costs, respiration rates, and cellular turnover demands.

1.1 Scope for Growth (SFG) and Energy Budgeting

A fish's total energy intake (C) is partitioned according to the standard bioenergetic balance equation:

$$C = (Gs + Gr) + (Mb + Ma + SDA) + (F + U)$$

C: Total ingested energy (Consumption).

Gs / Gr: Somatic growth and reproductive growth.

Mb / Ma / SDA: Standard metabolic rate (SMR), active metabolism, and specific dynamic action (heat increment of feeding).

F / U: Energy lost to feces (egestion) and nitrogenous waste (excretion).
Scope for Growth (SFG) represents the differential between total energy intake (C) and total metabolic loss (M). Maxima in somatic growth occur at the specific water temperature where SFG is optimized—where feed intake remains high while basal metabolic expenditure (Mb) leaves substantial excess energy for protein deposition.

The Thermal Performance Curve and Growth Zones
Every teleost species exhibits a distinct Thermal Performance Curve (TPC) that characterizes its growth response across three primary developmental zones:

2.1 Sub-Optimal Low Temperatures

Gastric Emptying & Digestion: Cold temperatures suppress digestive enzyme activity (e.g., pepsin, trypsin, lipase) and slow gastrointestinal motility. Gastric Emptying Time (GET) increases dramatically.

Appetite Suppression: Reduced neurological signaling via ghrelin and neuropeptide Y (NPY) depresses feed consumption.

Somatic Stagnation: While basal metabolic demands drop, energy intake drops faster, resulting in minimal growth or weight loss as energy stores are drawn down to maintain core cellular functions.

2.2 Thermal Optimum (Topt)
At the thermal optimum, the kinetic energy of metabolic enzymes aligns with optimal protein structure and lipid fluidity. Digestive enzymes operate at maximum efficacy, appetite peaks, and the Feed Conversion Ratio (FCR) drops to its minimum.

2.3 Sub-Optimal High Temperatures
Exponential SMR Acceleration: As temperatures surpass Topt, standard metabolic rate increases exponentially. The energetic cost of staying alive outweighs energy intake.

Dissolved Oxygen (DO) Inversion: Warm water exhibits lower physical solubility of dissolved oxygen. Simultaneously, the fish's tissue oxygen demand spikes due to accelerated tissue respiration. This disparity leads to systemic hypoxia.

Growth Collapse: Feed conversion efficiency plummets. Excess heat forces cellular energy to be diverted away from growth toward heat shock protein (HSP70, HSP90) expression, antioxidant defense, and respiratory ventilation.

The Ecological Perspective: Temperature-Size Rule (TSR)
In both wild populations and ecological research, temperature variations drive a developmental response known as the Temperature-Size Rule (TSR):

> **TEMPERATURE-SIZE RULE (TSR)**
>
> *Warmer Water Rearing ==> Higher Juvenile Growth Rate & Lower Maximum Adult Body Size*

Mechanistic Drivers of TSR:

1. Accelerated Cell Division vs. Cell Volume: Warm conditions accelerate cellular recruitment and developmental rate relative to cellular growth volume, causing rapid juvenile development but earlier maturation at smaller adult sizes.

2. Early Reproductive Axis Activation: High temperatures stimulate the hypothalamic-pituitary-gonadal (HPG) axis earlier in development, prompting an earlier bioenergetic transition from somatic muscle growth to gonadal development.

Comparative Thermal Trajectories across Key Cultured Species

Thermal response ranges differ significantly across functional groups (cold-water, temperate/eurythermal, and warm-water species):

Species Group	Model Species	Thermal Range (°C)	Topt for Growth (°C)	Primary Physiological Limitation at High Limits
Cold-Water	Atlantic Salmon (Salmo salar)	2 – 18	12 – 14	Oxygen transport limitation, cardiac failure, thermal stress
Temperate	Rainbow Trout (Oncorhynchus mykiss)	5 – 21	14 – 16	High SMR, lower mucosal immune barrier integrity
Eurythermal / Warm	African Catfish (Clarias gariepinus)	22 – 34	28 – 31	Slowed gastric clearance at <22°C, high waste toxicity at >34°C
Warm-Water	Nile Tilapia (Oreochromis niloticus)	20 – 36	28 – 32	Acute stress response, suppressed antibody synthesis at lower limits

Impacts of Diel Thermal Fluctuations

In natural water bodies and shallow aquaculture ponds, temperatures are rarely static; they exhibit diel temperature fluctuations (DTF) driven by day-night solar cycles.

Moderate Fluctuation (ΔT = ±2–4°C): Mild thermal cycling can stimulate growth rates (a phenomenon termed thermal compensation or growth hormesis) compared to rigid constant-temperature regimes. Enzyme expression shifts adaptively between day and night cycles.

Severe Fluctuation (ΔT > ±6°C): Rapid thermal shifts induce acute physiological stress. The energy spent continually adjusting cell membrane fluidity (homeoviscous adaptation) and resetting physiological pathways reduces energy available for somatic growth.

Strategic Implications for Aquaculture Bioenergetics

1. Precision Feed Management: Feeding regimes must be dynamically adjusted based on real-time water temperature. Feeding heavily during periods of extreme high or low water temperature causes feed waste, degrades water quality, and induces intestinal stress.

2. Dissolved Oxygen Management: Aeration capacity must scale non-linearly with rising water temperatures to satisfy the dual effect of lower oxygen solubility and heightened metabolic oxygen demand.

3. **Genetic and Breeding Strategies:** Selective breeding programs focusing on thermal tolerance markers (e.g., thermal performance breadth, expression of specific heat shock proteins) will be necessary to build resilience against climate-driven temperature fluctuations.

CONCLUSION

Temperature variation acts as an environmental regulator of fish growth, exercising control over metabolic rates, digestive kinetics, feed intake, and energy partitioning. While temperature increases up to a species' specific optimum enhance somatic growth by boosting feed conversion and enzyme kinetics, exceeding this limit causes rapid physiological degradation, metabolic exhaustion, and growth decline. Future research must continue exploring the epigenetics of thermal acclimation and metabolic gene regulation to better support sustainable aquaculture and marine conservation efforts in a warming world.

REFERENCES

1. Jobling, M. (1994). Fish Bioenergetics. Chapman & Hall, Fish and Fisheries Series 13, London.
2. Pörtner, H. O., & Farrell, A. P. (2008). Physiology and Climate Change. Science, 322(5902), 690–692.

3. Brett, J. R. (1979). Environmental factors and growth. Fish Physiology, 8, 599–675.
4. Sheridan, M. A. (1994). Regulation of lipid metabolism in fishes. Progress in Lipid Research, 33(3), 299–335.
5. Baroiller, J. F., et al. (2009). Temperature effects on sex determination and differentiation in fish. Journal of Fish Biology, 75(10), 2393–2441.
6. Van Der Kraak, G., & Pankhurst, N. W. (1997). Temperature effects on fish reproduction. Global Warming: Implications for Freshwater and Marine Fish, 159–176.