

Nernst's Partition Law in Solubility and Partitioning of Benzoic Acid in Aqueous-Organic Solvents System

Shiv Prakash Mishra

Assistant Professor in Chemistry (Guest), Faculty of Science, Dr. Ram Manohar Lohia Avadh University, Ayodhya-224001, (U.P.), India

ABSTRACT

Solubility and partition study of benzoic acid in two immiscible solvents like hydrophilic aqueous (water) and hydrophobic organic (benzene) layers have been reported by involving physico-analytical method. At room temperature (25°C), the solubility of solute benzoic acid in g/100g of solvents at different pH buffer solutions are analyzed by using titration method, where the partition or distribution coefficient of acid in these two solvents by shake-flask method. The solubility and distribution coefficient of benzoic acid in distilled water and in benzene-buffer solutions are determined at pH 4.0, 7.0 and 9.0 (acidic, neutral and basic). The analysis of benzoic acid solubility in g/100g of water solvent is being 0.142±0.033 and in different buffer solutions at pH 4.0, 7.0 and 9.0 it being 0.153±0.012, 0.186±0.145 and 0.148±0.708 respectively. Distribution coefficient for water-benzene solvent system is determined by introducing of Nernst partition law, $K = C_A/C_B$. The value of pH analysis on partition coefficient of benzoic acid for water-benzene solvent is being 0.636 and in buffer solutions as 0.841, 0.624 and 0.589 at pH 4.0, 7.0 and 9.0 respectively. Graph observation show that, the solubility of benzoic acid in buffer solutions as in acidic pH (4.0) is about slightly higher than basic (9.0) medium, but in comparing to distribution coefficient of acid in acidic medium is higher than both neutral (7.0) and basic pH.

Keywords: Benzoic acid, Solubility, Distribution coefficient, Nernst partition law equation, pH effect.

INTRODUCTION

Indeed, solubility and partition coefficient of solute in solvent was firstly performed by Berthelot and Jungfleisch¹, but it proved well by W. Nernst in 1891². Nernst studied the distribution of several solutes between two non-miscible or different appropriate pairs of solvents. Although, in literature about solubility and distribution coefficient of solute in solvents with its application³⁻⁷, partition of water-octanol⁸, and salvation thermodynamics in water-chloroform system have also been well reported⁹. The benzoic acid solubility and its dissociation process between water and benzene solvents have reported by titration¹⁰, or adapting of various physico-analytical methods¹¹⁻¹³. Concerning the study of solubility and partition coefficient of benzoic acid at 25°C as room temperature which having compatibility with Le Chatelier's principle¹⁴. In series of aromatic acids the benzoic acid (C₆H₅COOH) having poor solubility in cold water. In acid the carboxylic acid group is polarize during on dissociation partially in water by forming H-bonding and to produce benzoate anion (C₆H₅COO⁻) as well as a hydronium cation (H₃O⁺) in reaction. In saturated aqueous solution (aq), a little molar solubility of benzoic acid which has following equilibrium-



The dissociation equilibrium of benzoic acid is expressed as following-

$$Kc = \frac{[\text{H}^+][\text{C}_6\text{H}_5\text{COO}^-]}{[\text{C}_6\text{H}_5\text{COOH}(aq)]}$$

Here, the Kc is dissociation constant of benzoic acid. Knowing, the benzoic acid is a colorless crystalline solid substance having M.F. C₆H₅COOH and M.P. 121-122°C with pleasant smell¹⁵, and in early 20th century its salt has been used in industrial and medicinal purposes^{16,17}. In present study we have been selected a benzoic acid because its suite solubility in variety of polar and non-polar solvents such as in water (H₂O) benzene (C₆H₆), carbon tetrachloride (CCl₄), chloroform (CHCl₃), alcohol (C₂H₅OH), acetone (CH₃COCH₃) and in liquor ammonia (liq.NH₃), etc. The benzoic acid solubility in

water and in benzene with buffer solutions of pH acidic (4.0), neutral (7.0) and basic (9.0) medium are determined by using titration method and partitioning study of that acid in both liquid solvents by shake-flask method. The solute benzoic acid concentration in water and benzene solution is analyzed by applying acid-base titration method.

In article, we are reported a Nernst's partition law in solubility and partitioning of benzoic acid between two solvents like as hydrophilic aqueous (water) and hydrophobic organic (benzene) layer which are determined by using titration method. In this work we are using equation derived from Nernst distribution coefficient law, which have used and improved to finding an accurate results with value of K . Mathematically, the expression of Nernst distribution equation is given as-

$$K = C_A / C_B \dots\dots\dots (1)$$

Here, the C_A and C_B are concentration of substances as for solvent A and B, and K is a constant and called Nernst's partition coefficient or distribution coefficient at fixed temperature, where its value is independent of actual concentration of the solutions. When a solute is shaken with two immiscible solvents, at equilibrium both solvents are saturated with the solute. Since the solubility also represents concentration, thus we can write the Nernst's distribution law as in following equation-

$$K = C_A / C_B = S_A / S_B \dots\dots\dots (2)$$

where S_A and S_B are the solubility of solute in the two solvents A and B.

In case when solute undergoing partition has molecular weight in one solvent say (organic solution) n times higher than water at equilibrium, the distribution ratio K becomes-

$$K = C(aq) / C(org)^{1/n} \dots\dots\dots (3)$$

Here, $C(aq)$ is equilibrium concentration of solute (benzoic acid) in aqueous solution (water), and $C(org)$ is equilibrium concentration of solute in organic solvent (benzene). Since, benzoic acid exists as dimer in aprotic solvents like in benzene (i.e., $n=2,3$)¹⁸, hence the modified Nernst's distribution law is valid-

$$K = C(aq) / n\sqrt{C(org)} \dots\dots\dots (4)$$

where, n is known as degree of association with K as partition ratio or coefficient.

EXPERIMENTAL

(1)- Materials: In experimental procedure, the all required chemicals and solvents which are used in this study as an analytical reagent grade on laboratory based as well as the solvent distilled water is also freshly prepared which is used during throughout the work. The carbonate free sodium hydroxide (NaOH) solutions are prepared as work suggested by Vogel¹⁹. A Research Lab Fine Chemicals Industries, Mumbai, have provided solid benzoic acid, (C_6H_5COOH) and the Merck Specialties (Pvt.) Limited, Mumbai, India, to buffer solutions as of pH 4.0, 7.0 and 9.0 value. In this work those calibrated volumetric glassware used which is an 'A' class standard.

(2)- Methodology:

(a)- The solubility (S) determination of benzoic acid: At 25°C temperature we are applying the acid-base titration method, for estimation of benzoic acid solubility at varies pH buffer solution. Now, in different beakers take 100ml buffer solution of pH 4.0, pH 7.0 and pH 9.0, and then in every beaker we added solid benzoic acid about 200mg, and for producing saturated solution it stirred with glass rod. Notably, some solid un-dissolved and must be left. If required these solution is heated. At 25°C, cooling these solution which are prepared and into dry conical flask (say W_1) it withdrawn a 5.0ml of these solution. Again weight (say W_2) of 5.0ml solution contained conical flask and indicator phenolphthalein is used and it titrate against to these solution with prepared of 0.05N NaOH solution. Records the burette reading (V) as appearing pink colour show it end point. As given below described formula we determine the benzoic acid solubility (S) in gram/100gram of solvents and to compare with water (distilled) as blank against solubility²⁰. Here, the W_4 ($0.120 \times 0.05N \times V$) is as g/w of solute substance $\times N$ of solution $\times ml$ of solution used.

If, conical flask weight (empty) and with solution (5.0ml) are = W_1 and W_2 , respectively

then, the solution weight ($W_2 - W_1$) = W_3

and, if the solute weight (C_6H_5COOH) = W_4

then, the solvent weight, ($W_3 - W_4$) = W_5

Thus, solubility (S) = solute weight (W_4) $\times 100$ / solvent weight = (W_5)

(b)- The partition coefficient (K) determination of benzoic acid: By the using acid-base and shake-flask method we estimated partition coefficient (K) of benzoic acid at different pH. Where, in a beaker we prepared 10 % benzoic acid solution in benzene (say 2B solution). Here, the four different type of solutions are prepared in separating funnel, where

buffer solution having the same pH (4.0) for each all sample, is given below-

- (1). Buffer solution (40 ml) + 2B solution (40 ml)
- (2). Buffer solution (40 ml) + benzene (10 ml) + 2B solution (30 ml)
- (3). Buffer solution (40 ml) + benzene (15 ml) + 2B solution (25 ml)
- (4). Buffer solution (40 ml) + benzene (20 ml) + 2B solution (20 ml)

At room temperature, we used a water-bath incubator shaker for where all prepared solutions with in flasks have been shaken for one day (24hours) and allowing to stand it about half hour to get equilibrium. After then we obtained that the used separating funnel or flask is containing lower aqueous with upper as benzene layer. In dry beaker, firstly we removed the lower layer (aqueous) from each flask by retaining of upper layer (benzene) within separating flask. Now, in a dry conical flask we pipette out aqueous layer of 10 ml, and against with 0.01N NaOH solution we titrated it, by using indicator as phenolphthalein. The pink color as it end point. We record this reading well. Benzene layer (5.0 ml) is pipette out in an another using dry conical flask with adding a 10 ml of distilled water. Using of indicator as phenolphthalein also, these prepared solution with 0.1N NaOH solution against titrated. A pink color is appearance as it end point of reaction and record this observe burette reading. Notably, we have followed for buffer solution of pH 7.0 and pH 9.0, the same above described procedure with to blank sample as distilled water. By applying normality and modify Nernst partition formula for aqueous water and organic benzene phase or layer, we are estimate as well as to calculate the partitioning (*K*) behavior of solute in solvent (in moles/litre). The following formula have been given below-

(1)- Benzoic acid (solute) concentration for aqueous (water) layer :

As determination of normality it is calculated by applying following normality equation-

$$N_1V_1 = N_2V_2 \dots \dots \dots (i)$$

Here, the N_1 is aqueous layer normality written as, $N(aq)$ and the $N(aq) = 0.01 \quad V_2 / 10 \dots \dots (ii)$

and, the N_2 is normality of base NaOH which involve for titration as $N(org) = 0.01 \text{ N}$,

thus, the $N(aq) = C(aq) \dots \dots \dots (iii)$

The V_1 is taken volume of aqueous layer (10 ml), and V_2 the consumed as burette reading of volume of base NaOH.

(2)- Benzoic acid (solute) concentration for organic (benzene) layer :

As determination of normality it is calculated by applying following normality equation-

$$N_3V_3 = N_4V_4 \dots \dots \dots (iv)$$

Here, the N_3 is organic layer normality as $N(org), = 0.1 \quad V_4 / 5 \dots \dots (v)$

and the N_4 is normality of base NaOH which used for titration as $N(org) = 0.1 \text{ N}$, thus the,

$$N(org) = C(org) \dots \dots \dots (vi)$$

The V_3 is taken volume of organic layer (5.0 ml), and V_4 as the burette reading (consumed) of volume of base NaOH.

(3)- Partition coefficient (K) of solute for aqueous and organic solvents system:

The partition coefficient of aqueous-organic solvents system which is determined from applying equation-

$$K = \sqrt{C(org) / C(aq)}, \text{ or } = C(aq) / C(org)^{1/2} \dots \dots (vii)$$

Thus, at equilibrium, $K = (C_w / C_o)^{1/2}$

Where, C_w is stand for concentration of solute (benzoic acid) in aqueous water layer and C_o for concentration of solute in organic layer.

RESULT AND DISCUSSION

The solubility and distribution of solute is depends on various factors which are as temperature, pressure, intermolecular forces, and nature of solute-solvent with solute dissociation in solvent as in ionic form and H-bonding *etc.* The addition of salts in water gives rise a slight changing because solute-solute and solute-solvent interaction²¹. The solute benzoic acid in aqueous solution has less solubility and having weaker electrolyte behaviour²². From table-1 we have reported that the solute benzoic acid's solubility in gram/100gram of solvent, at room temperature about 25°C, as in water (as distilled) is being to 0.142 ± 0.033 and at three pH 4.0, pH 7.0 and pH 9.0 it is varies to 0.153 ± 0.012 , 0.186 ± 0.145 and 0.148 ± 0.708 g/100g with solvent for benzene-buffer solution, respectively. This observation reveals that the value is being maximum at

neutral pH (7.0) for benzene and buffer solution. The figure-1 have shown benzoic acid solubility with pH effect, where graph is plotting between solubility of solute in g/100g of solvent versus varied pH of buffer solutions and water. These suggested that the acid solubility as in acidic range is becoming higher slightly in comparison of basic medium due to their strength of ionic dissociation.

Table-1. At 25°C, The Solubility Analysis Of Benzoic Acid

S.N.	Using Solvents	pH	Mean Solubility (S) (in g/100g of solvent)
1-	Water (distilled)	-	0.142 ± 0.033
2-	Benzene-buffer solution	4.0	0.153 ± 0.012
3-	Benzene-buffer solution	7.0	0.186 ± 0.145
4-	Benzene-buffer solution	9.0	0.148 ± 0.708




Figure-1. The pH effect on benzoic acid (solute) solubility

As from table-2, we have reported the pH behavior analysis of partitioning of benzoic acid in phase of water and benzene with acidic, neutral and basic buffer solutions. Where, the value for benzene-water phase it is being 0.636 as well as the combine value as 0.841, 0.624 and 0.589 is being for benzene-buffer solutions at different range pH 4.0, 7.0 and 9.0, respectively. The figure-2 has been shown, where, the graph is plotted between partitioning (K) and the buffer solutions of different pH value. Graph observation indicate that, the benzoic acid distribution coefficient in acidic pH medium is higher than in comparison of that the neutral as well as in basic medium both.

Table-2. pH analysis on partition coefficient of Benzoic Acid

S.N.	Using Solvents	pH	Partition coefficient (K)
1-	Benzene-water solvent	-	0.636
2-	Benzene-buffer solution	pH 4.0	0.841
3-	Benzene-buffer solution	pH 7.0	0.624
4-	Benzene-buffer solution	pH 9.0	0.589




Figure-2. The pH effect on partition coefficient of benzoic acid

Graphically, for aqueous-organic solvent system, the both graphs are plotted in respect of solubility (S) and partitioning or distribution coefficient (K) of solute in solvents with different pH containing buffer solutions and effect of pH on solubility and partitioning of benzoic acid. For distribution of solute in aqueous-organic solvents, the study reveals that as C_w and $C_o^{1/n}$ (concentration of water and organic layer) ratio is remain constant, whereas, the C_w and C_o ratio may not remain constant. It indicates the formation of H-bonding by association of molecules in dimerization phenomenon between molecules of benzoic acid in organic layer and in aqueous layer the remaining monomer molecules. In water the benzoic acid solubility is directly proportional with temperature, but at higher temperature than 30°C , the acid behaves differently. The capability of benzoic acid to dissociate which leads to reduce the value of K^{10} , with inductive effect inside the acid molecule resulting polarization of bond ²³.

CONCLUSION

In conclusion, we have reported the Nernst's partition law in study of solubilization and partitioning of solute benzoic acid in immiscible aqueous water and organic benzene solvent layers by involving physico-analytical method. Solubility of benzoic acid in different pH buffer solutions are analyzed by using titration method, and partition or distribution coefficient of such acid in these two solvents by shake-flask method at room temperature (25°C). Where, the solubility and distribution coefficient of acid in distilled water and in benzene-buffer solutions are determined at pH 4.0, 7.0 and 9.0 (in acidic, neutral and basic medium). Benzoic acid solubility analysis in g of solute/100 g of solvent water is being 0.142 ± 0.033 and in different buffer solutions at pH 4.0, 7.0 and 9.0 it is 0.153 ± 0.012 , 0.186 ± 0.145 and 0.148 ± 0.708 respectively. The distribution coefficient for water-benzene solvent system is determined by introducing of Nernst partition law, $K = C_A/C_B$. For pH analysis on partition coefficient of benzoic acid in water-benzene solvent is being 0.636 and in buffer solutions it as 0.841, 0.624 and 0.589 at pH 4.0, 7.0 and 9.0 respectively. The graph observation show that solubility of benzoic acid in buffer solutions at acidic pH (4.0) is about slightly higher than basic (9.0) medium, but in comparing to distribution coefficient of acid in acidic medium is higher than other neutral (7.0) and basic pH.

ACKNOWLEDGEMENTS

These work supported by various respective Journals and books. The author thanks to gratefully acknowledge support of Faculty of Science, Dr. Ram Manohar Lohia Avadh University, Ayodhya-224001, (U.P.), India, for providing help in useful discussion and necessary facilities to valuable data.

REFERENCES

- [1]. M. Berthelot and J. Jungfleisch, *Am. Chim et Phys.*, 26, 396, 1872.
- [2]. W. Nernst, *Z. Phys. Chem.*, 8, 110, 1891.
- [3]. A. Leo, C. Hansch and D. Elkins, *Chem. Rev.*, 71, (6), 525-616, 1971.
- [4]. E. S. Machlin, *An Introduction to Aspect of Thermodynamics and Kinetics Relevant to Materials Science*, 3rd Edition, Secondary, Elsevier, Amsterdam, 98, 2007.
- [5]. R. Stallman and A. Ngan, *Solidification and Modern Physical Metallurgy*, 8th Edition, Elsevier-Butterworth-Heinemann, Amsterdam, 93-120, 2014.

- [6]. A. Berthod and S. Carda-Broch, *J. Chromatography A*, 1037, 3-14, 2004; A. Berthod, and N. Mekaoui, *J. Chromatography A*, 1218, (36), 6024-6030, 2011.
- [7]. N. N. Hidayah, S. Z. Abidin, *Minerals Engineering*, 112, 103-111, 2017.
- [8]. H. Cumming and C. Rucker, *ACS Omega*, 9, 6244-6249, 2017.
- [9]. J. Kraml, F. Hofer, A.S. Kamenik, F. Waibl, U. Kahler, M. Schauerperl and K. R. Liedl, *J. Chem. Inf. Model.*, 60, (8), 3843-3853, 2020.
- [10]. S. J. Khouri, *American J. Analytical Chem.*, 6, 429-436, 2015.
- [11]. Frederick T. Wall, *J. Am. Chem. Soc.*, 64, (2), 472-473, 1942.
- [12]. K. Sagarik, S. Chaiwongwattana and P. Sisot, *Chemical Physics*, 306, (1-3), 1-12, 2004.
- [13]. X. Shi, L. Qiao, and G. Xu, *J. Chromatography A*, 1420, 1-15, 2015.
- [14]. P. Atkins and J. de Paula, *Physical Chemistry*, 9th Edition, W. H. Freeman and Company, New York, 2010.
- [15]. I. L. Finar, *Organic Chemistry*, Vol-1, 6th Edition, Dorling Kindersley (India) Pvt. Ltd., 2007.
- [16]. C. C. Xu, B. Wang, Y. Q. Pu, J. S. Tao, and T. Zhang, *Chin. J. Nat. Med.*, 15, (10), 721-731, 2017.
- [17]. S. Shi, D. Fan, H. Xiang and H. Li, *Food Chem.*, 237, 198-204, 2017.
- [18]. L. E. Strong, C. L. Brummel, R. Ryther, J. R. Radford and A. D. Pethyridge, *J. Solution Chem.*, 17, 1145-1167, 1988.
- [19]. A. Vogel, *Text book of Quantitative Chemical Analysis*, 5th Edition, Longman, Harlow, 1989.
- [20]. S. P. Mishra, *Chemical Sci. Int. J.*, 30, (3), 40-45, 2021.
- [21]. V. F. Sergeeva, *Russian Chemical Reviews*, 34, 309-318, 1965.
- [22]. A. Albert and E.P. Serjeant, *The Determination of Ionisation Constants*, 3rd Edition, Chapman and Hall, London, 1984.
- [23]. G. T. Solomons and C. B. Fryhle, *Organic Chemistry*, 10th Edition, John Wiley & Sons, Hoboken, 2011.